

## Effect of Zinc Ion on Formation of the Fertilization Membrane in Sea Urchin Eggs

SHOGO NAKAMURA, CHIKA OHMI and MANABU K. KOJIMA

*Department of Biology, Faculty of Science,  
Toyama University, Toyama 930, Japan*

**ABSTRACT**—The effect of  $Zn^{2+}$  on the formation of the fertilization membrane was observed in eggs of the sea urchin, *Anthocidaris crassispina*. When the eggs were transferred into sea water containing more than  $5 \mu M Zn^{2+}$  within 10 sec after insemination, formation of the fertilization membrane was inhibited. This inhibition was irreversible. Although the time of the first cleavage was delayed in a dose-dependent fashion, cleavage occurred normally in the presence of up to  $1 mM Zn^{2+}$ . Electron microscopic examinations revealed that the exocytosis of cortical granules progressed. However, the elevation and the hardening of the vitelline layer were blocked by more  $5 \mu M Zn^{2+}$ .

### INTRODUCTION

When sea urchin eggs are fertilized or artificially activated, cortical granules, which are located beneath the plasma membrane, are exocytotically discharged and cortical granule components are released into the perivitelline space. Some of the components hydrate and elevate the fertilization membrane, while others are incorporated into the fertilization membrane to make it thick and tough. Thus the fertilization membrane is formed around the eggs [see references for review 1, 6].

It is well known that  $Zn^{2+}$  inhibits the formation of the fertilization membrane [2, 4, 7]. However, the mechanism of the inhibition by  $Zn^{2+}$  is unknown. Moreover, not much is known about the mechanism of the fertilization membrane formation itself. We consider that clarifying the inhibitory mechanism by  $Zn^{2+}$  may provide information concerning the mechanism of formation of the fertilization membrane. As our first step, we observed in detail the effect of  $Zn^{2+}$  on formation of the fertilization membrane. The results suggest that  $Zn^{2+}$  inhibits the elevation and hardening of the fertilization membrane occurring after the cortical reaction in the process of the fertilization membrane formation.

### MATERIALS AND METHODS

Gametes from the sea urchin, *Anthocidaris crassispina*, were collected by intracoelomic stimulation with  $0.5 M KCl$ . Sperm was kept 'dry' in a refrigerator until use; eggs were spawned into filtered sea water, and washed three times by sedimentation. Experiments were done at  $25-29^{\circ}C$ .

Small amounts of  $Zn^{2+}$  from a stock solution ( $100 mM ZnCl_2$ ) in deionized water were added to the experimental sea water. The eggs were transferred to the sea water containing  $Zn^{2+}$  10-60 sec after insemination.

Formation of the fertilization membrane was observed with an inverted microscope (Olympus CK-2). Micrographs were taken on Fuji Neopan F film.

Eggs for electron microscopy were fixed in 2% glutaraldehyde in 80% sea water (pH 7.0), and post-fixed in 1%  $OsO_4$  in 50 mM sodium cacodylate (pH 7.2). After dehydration through an ethanol series, they were infiltrated and embedded in epoxy resin (Epok 812, Oken Shoji Co., Ltd., Tokyo). Ultrathin sections were stained with uranyl acetate and Reynold's lead citrate and examined with a JEOL 100C electron microscope operated at 80 kV.

Accepted June 30, 1988

Received April 5, 1988

## RESULTS AND DISCUSSION

The eggs of *Anthodiaris crassispina* were treated with various concentrations of  $Zn^{2+}$  from 10, 30 and 60 sec after insemination. Figure 1 shows the




Fig. 1. Effect of  $Zn^{2+}$  on formation of the fertilization membrane in *Anthodiaris crassispina* eggs. At 10 sec (○), 30 sec (●), or 60 sec (△) after insemination, the eggs were transferred into sea water containing various concentrations of  $Zn^{2+}$ .

effects of increasing  $Zn^{2+}$ -concentration on formation of the fertilization membrane. Although the rate of inhibition varied from experiment to experiment, formation of the fertilization membrane was completely prevented when more than  $5 \mu M$   $Zn^{2+}$  was applied within 10 sec after insemination (Figs. 1 and 2b). This inhibition was irreversible. When eggs without a fertilization membrane in  $5 \mu M$   $Zn^{2+}$ -containing sea water were removed and placed in fresh sea water, they could not form a fertilization membrane. The surfaces of the eggs in  $Zn^{2+}$ -containing sea water were rougher than those of unfertilized eggs in normal sea water (Fig. 2b, c).

To confirm that fertilization occurred in the eggs without a fertilization membrane in  $Zn^{2+}$ -containing sea water, we measured percentage of cleavage in those eggs (Fig. 3). The time of the first cleavage was delayed by  $Zn^{2+}$  in a dose-dependent manner, but cleavage occurred normal-

dependent manner, but cleavage occurred normally in the presence of up to  $1 mM$   $Zn^{2+}$ . For




Fig. 2. Formation of the fertilization membrane in eggs treated or untreated with  $Zn^{2+}$ . (a) Control, (b)  $5 \mu M$ , and (c)  $1 mM$   $Zn^{2+}$ . Bar,  $50 \mu m$ .







Fig. 3. Percentage of cleaved eggs in the presence of various concentrations of  $Zn^{2+}$ . At 10 sec (○), 30 sec (●), or 60 sec (△) after insemination, the eggs were transferred into sea water containing  $Zn^{2+}$ . The percentage was measured at 55 min (solid line) and 65 min (dotted line) after insemination.

Fig. 4. Electron micrographs of sea urchin eggs. The eggs were fixed at 10 min after insemination. An unfertilized egg (a), a fertilized egg (b), and a fertilized egg treated with  $5 \mu M$   $Zn^{2+}$  from 10 sec after insemination (c). CG, cortical granule; FM, fertilization membrane; MV, microvillus; YG, yolk granule; UC, undispersed contents of cortical granule. Bar,  $1 \mu m$ .






FIG. 5. High-magnification views of portions of the eggs shown in Fig. 4 (a) and (c). (a) Unfertilized egg, and (b)  $Z^{2+}$ -treated egg. CG, cortical granule; VL, vitelline layer; PM, plasma membrane; YG, yolk granule; UC, undispersed contents of cortical granule. Bar,  $0.5 \mu\text{m}$ .

ly in the presence of up to 1 mM  $Zn^{2+}$ . For example, 3% of eggs transferred into 1 mM  $Zn^{2+}$ -containing sea water within 10 sec after insemination cleaved by 55 min after insemination, but the rate of cleavage had recovered to 82% by 65 min after insemination. This result shows that fertilization occurred normally in  $Zn^{2+}$ -treated eggs.

Next, we examined by electron microscopy whether the cortical granules react normally in eggs treated with  $Zn^{2+}$  (Figs. 4 and 5). Exocytosis of cortical granules occurred in  $Zn^{2+}$ -containing sea water (Figs. 4c and 5b), i.e., the membrane of cortical granules fused with the plasma membrane of the egg and the contents were released out of the egg. However, dispersion of the contents was incomplete, and part of them were undispersed by 10 min after insemination. Moreover, detachment of the vitelline layer from the plasma membrane and hardening of the vitelline layer were blocked by more than 5  $\mu$ M  $Zn^{2+}$ . This resulted in rupture of the vitelline layer by the pressure from influx of water into the perivitelline space or swelling of the cortical granule components. The undispersed contents may relate to the detachment and hardening of the vitelline layer.

In normal formation of the fertilization membrane, the following processes occur [1, 6]. (a) Intracellular concentration of free  $Ca^{2+}$  is increased by fertilization or artificial activation. (b) This  $Ca^{2+}$  triggers the cortical granule exocytosis. (c) Proteases in the cortical granules break the attachment between the vitelline layer and the plasma membrane of the eggs. (d) The vitelline layer is elevated by hydration and/or osmotic effects resulting from the secretion of cortical granule contents. (e) The vitelline layer is hardened through cross-linkage and structuralization with cortical granule-derived proteins, such as ovoperoxidase [1, 6] and proteoliasin [8]. The results we obtained in this study suggest that  $Zn^{2+}$  inhibits steps (c), (d) and (e) in the process of formation of the fertilization membrane.

$Zn^{2+}$  is a potent animalizing agent in sea urchin embryos. However, we do not know how animalization is induced by  $Zn^{2+}$ . Lallier [3] suggested that when  $Zn^{2+}$  binds to proteins, their structure and biochemical properties may become altered, so that abnormal development occurs. In fertiliza-

tion membrane formation,  $Zn^{2+}$  may exert the same effect on proteins. We are now searching  $Zn^{2+}$ -binding protein in the cortical granule components of unfertilized eggs. In addition,  $Zn^{2+}$  has an affinity for -SH groups, and action of  $Zn^{2+}$  on the enzymes whose activity depends on -SH groups has also been suggested [5]. It should be noted that -SH groups may play an important role in the elevation and the hardening of the vitelline layer.

#### ACKNOWLEDGMENTS

We are grateful to Dr. K. Aketa (Hokkaido University) for his critical reading of this manuscript. We also thank Drs. R. Kamiya, S. Higashi-Fujime and S. Asakura (Nagoya University) for their help in electron microscopy.

#### REFERENCES

- 1 Kay, E. S. and Shapiro, B. M. (1985) The formation of the fertilization membrane of the sea urchin egg. In "Biology of Fertilization, Vol. 3". Ed. by C. B. Metz, and A. Monroy, Academic Press, New York, pp. 45-80.
- 2 Kobayashi, N. (1971) Fertilized sea urchin eggs as an indicatory material for marine pollution bioassay, preliminary experiments. *Publ. Seto Mar. Biol. Lab.*, **18**: 379-406.
- 3 Lallier, R. (1975) Animalization and vegetalization. In "The Sea Urchin Embryo: Biochemistry and Morphogenesis." Ed. by G. Czihak, Springer-Verlag/Berlin/Heidelberg/New York, pp. 473-509.
- 4 Murakami, H., Hayakawa, M., Fujii, T., Hara, T., Itami, Y., Kishida, A. and Nishida, I. (1976) The effects of heavy metals on developing sea urchin eggs. *Okayama Igakkai Zasshi*, **88**: 39-50. (in Japanese)
- 5 Rulon, O. (1955) Developmental modifications in the sand dollar caused by zinc chloride and prevented by glutathione. *Biol. Bull.*, **109**: 316-327.
- 6 Schuel, H. (1985) Functions of egg cortical granules. In "Biology of Fertilization, Vol. 3". Ed. by C. B. Metz, and A. Monroy, Academic Press, New York, pp. 1-43.
- 7 Sugiyama, M. (1949) Studies on the artificial parthenogenesis of the sea urchin eggs. VIII. Effect of heavy metal ions on membrane formation. *Zool. Mag. (Tokyo)*, **58**: 137-138. (in Japanese)
- 8 Weidman, P. J., Kay, E. S. and Shapiro, B. M. (1985) Assembly of the sea urchin fertilization membrane: Isolation of proteoliasin, a calcium-dependent ovoperoxidase binding protein. *J. Cell Biol.*, **100**: 938-946.